Contents

Summary of Major Findings	
Precipitation	
Ground Water	
Surface Water	2
Water Quality	2
Hydrologic Budgets	3
Introduction	
Purpose and Scope	
Description of Study and Study Area	
Overview of Study	4
Description of Study Area	8
Hydrologic Processes and Framework	10
Hydrologic Processes	
Local Hydrologic Framework	
Climate	
Geology	
Ground Water	
Surface Water	
Regional Ground-Water Framework	
-	
Water Resources	
Ground-Water Resources	
Characteristics of Major Aquifers	
Overview of Other Aquifers	
Ground-Water Levels	
Temporal Trends	
Comparisons between Madison and Minnelusa Aquifers	
Comparisons for Other Aquifers	58
Responses to Climatic Conditions	
Ground-Water Quality	
Background	
General Characteristics for Major Aquifers	
General Characteristics for Minor Aquifers	
Susceptibility to Contamination	
Ground-Water Quality Relative to Water Use	
Surface-Water Resources	
Streamflow Characteristics	
Streamflow Variability	
Annual Yield Characteristics	
Surface-Water Quality	
Standards and Criteria	
General Characteristics	
Human Influences on Water Quality	
Surface-Water Quality Relative to Water Use	92
Hydrologic Budgets	94
Methods for Estimating Basin Yield and Recharge	
Ground-Water Budgets	
Surface-Water Budgets	
Combined Ground-Water and Surface-Water Budget	
Madison and Minnelusa Flow System	
Flowpaths	
Springs	108
Future Information Needs	113
References	114
Definition of Torms	110

Figures—Continued

66-70.	Hydrographs illustrating:	
00-70.		EC
	66. Temporal trends in ground-water levels	56
	67. General similarities in water levels for some colocated Madison/Minnelusa wells with	
	confined conditions	
	68. Large hydraulic separation for colocated Madison/Minnelusa wells with unconfined conditions	
	69. Generally separated water levels for some colocated Madison/Minnelusa wells	58
	70. Colocated Minnelusa/Minnekahta and Deadwood/Madison wells	58
71-77.	Maps showing:	
	71. Water temperature in the Madison aquifer	63
	72. Specific conductance in the Madison aquifer	
	·	
	73. Specific conductance in the Minnelusa aquifer	
	74. Specific conductance in the Inyan Kara aquifer	
	75. Hardness in the Inyan Kara aquifer	67
	76. Sulfate concentrations in the Minnelusa aquifer	68
	77. Radon concentrations in the Deadwood aquifer	
78.	Boxplots of concentrations of nitrite plus nitrate for selected aquifers	70
79.	Photographs of potential sources of nitrogen in ground water	
80.	Graphs showing duration curves of daily mean streamflow	73
81.	Graphs showing variations in mean monthly streamflow for basins representative of	
	hydrogeologic settings	74
82.	Map showing basin yields for selected streamflow-gaging stations	75
83.	Map showing comparison between surface-drainage areas and contributing ground-water	
	areas for streamflow-gaging stations in Limestone Plateau area	76
84.	Schematic diagram illustrating recharge and streamflow characteristics for selected outcrop types	
85.	Map showing generalized average annual yield efficiency, water years 1950-98	
86.	Map showing estimated annual yield potential for the Black Hills area, water years 1950-98	
87.	Graph showing relations between dissolved solids and specific conductance by hydrogeologic setting	82
88.	Graph showing relations between specific conductance and streamflow for a selected	
	site within each hydrogeologic setting	83
89.	Diagrams of median concentrations of common ions by hydrogeologic setting	
90.	Map showing median sulfate concentrations in surface water	
91.	Map showing median uranium concentrations in surface water	
92.	Photographs of activities that can impact surface-water quality	
93.	Map showing maximum selenium concentrations in surface water	
94.	Graph showing downstream progression of pH for selected streams influenced by acid-mine drainage	88
95.	Photographs of an abandoned mine, acid-mine drainage, and a bog-iron area	
96.	Graph showing changes in sulfate and sodium concentrations at Bear Butte Creek near Deadwood	
97.	Graph showing comparison of dissolved copper concentrations to hardness-adjusted	00
37.		00
00	chronic and acute aquatic criteria for Bear Butte Creek near Deadwood	
98.	Map showing maximum arsenic concentrations in surface water	
99.	Graph showing nitrite plus nitrate concentrations in Annie Creek near Lead, 1988-97	91
100.	Boxplots of concentrations of dissolved nitrite plus nitrate by hydrogeologic setting,	
	with Annie Creek separated from other crystalline core sites	91
101.	Boxplots of concentrations of dissolved nitrite plus nitrate within the Rapid Creek Basin	
102.	Pie chart showing evapotranspiration, runoff, and precipitation recharge as percentages	-
102.	of annual precipitation	94
100		
103.	Schematic showing components considered for ground-water budgets	
104.	Photograph of headwater springflow in the Limestone Plateau area	
105.	Photograph of artesian springflow at Evans Plunge	96
106.	Pie charts showing percentages of average annual budget components for bedrock	
	aguifers for water years 1950-98	97
107.	Pie charts showing percentages of average annual budget components for the Madison	
107.	and Minnelusa aquifers in South Dakota and Wyoming for water years 1950-98	98
100	· · · · · · · · · · · · · · · · · · ·	30
108.	Pie chart showing percentages of water use from the Madison and Minnelusa aquifers,	00
	by category, for South Dakota counties in the Black Hills area for water years 1987-96	99
109.	Graph showing annual recharge to the Madison and Minnelusa aquifers, in the Black Hills	
	of South Dakota and Wyoming, water years 1931-98	99
110.	Photographs of the Cheyenne and Belle Fourche Rivers	100
111.	Schematic diagram showing average hydrologic budget components for the study area,	
	water years 1950-98	101
112.	,	101
112.	Schematic diagram showing generalized average streamflow relative to surface geology	101
	and depletions, water years 1950-98	
113.	Schematic diagram showing fractionation of stable oxygen isotopes	102
114.	Map showing generalized flowpaths in the Madison aquifer in the Black Hills area of	
	South Dakota and Wyoming	103
115.	Graph showing estimated tritium concentrations in precipitation for the Black Hills area	
110.	and decay curves for selected years	104
110	· · · · · · · · · · · · · · · · · · ·	104
116.	Boxplots of tritium concentrations for selected ground-water and surface-water	40.
	samples collected during 1990-98 in the Black Hills area	104
117.	Map showing tritium occurrence for selected sample sites in the Black Hills area	
118.	Map showing concentrations of δ^{18} 0 in Madison and Minnelusa aquifers in the Rapid City area	
119.	Photographs of dye testing along Boxelder Creek	
120.	Graph showing long-term streamflow and precipitation trends for gaging station	
0.	06409000, Castle Creek above Deerfield Reservoir	108
101	Graph showing long-term streamflow trends for gaging station 06402000, Fall River at Hot Springs	
121.		
122.	Photograph of red sediment at Cascade Springs	110
123.	Schematic diagrams illustrating the development and abandonment of an artesian spring	
	at the Mammoth Site of Hot Springs	111

Other Abbreviations, Acronyms, and Symbols Used

milligrams per liter mg/L micrograms per liter μg/L mL milliliters μS/cm microsiemens per centimeter at 25 degrees Celsius pCi/L picocuries per liter per mil ‰ TU tritium units less than < greater than > \geq greater than or equal to approximately equal to isotopic ratio of oxygen-18 (¹⁸0) to oxygen-16 (¹⁶0) δ^{18} 0 **DENR** South Dakota Department of Environment and Natural Resources GIS Geographic information system GWSI Ground Water Site Inventory database **USEPA** U.S. Environmental Protection Agency MCL Maximum Contaminant Level MSL Mean sea level **SMCL** Secondary Maximum Contaminant Level U.S. Geological Survey USGS Boxplots are a useful and concise graphical display for summarizing the distribution of a data set. Two different types of boxplots are used in this report. In both types, the center of the data (known as the median) is shown as the center line of the box. The variation or spread of the data (known as the interquartile range) is shown by the box height. Maximum The first type is a truncated boxplot, and is used for all 90th percentile boxplots that do not show water-quality data. In the 75th percentile truncated boxplot, the whiskers are drawn only to the Median 90th and 10th percentiles of the data set. Thus, values 25th percentile included in largest 10 percent and the smallest 10th percentile 10 percent of the data are not shown. The maximum Minimum and minimum values for the data set are shown. Outlier data value more than 3 times the The second type is a standard boxplot, and is used for all interquartile range outside the quartile boxplots that show water-quality data. In the standard Outlier data value less than or equal to 3 and boxplot, the whiskers are drawn only to the last data more than 1.5 times the interquartile range value that is within 1.5 times the interquartile range outside the quartile (height of the box). Values outside 1.5 times the Data value less than or equal to 1.5 times interquartile range are called "outliers." For waterthe interquartile range outside the quartile quality data, these outliers are of interest when 75th percentile comparing to water-quality standards and general Median distribution of extreme values. 25th percentile Data value less than or equal to 1.5 times the

<u>▼</u> Water table

9

Spring

interquartile range outside the quartile